

The large sieve with sparse sets of moduli

Stephan Baier

22.08.05

Address of the author:

Stephan Baier
Jeffery Hall
Department of Mathematics and Statistics
Queen's University
University Ave
Kingston, Ontario, Canada
K7L 3N6

e-mail: sbaier@mast.queensu.ca

Abstract: Extending a method of D. Wolke [7], we establish a general result on the large sieve with sparse sets $\mathcal{S}$ of moduli which are in a sense well-distributed in arithmetic progressions. We then apply our result to the case when $\mathcal{S}$ consists of squares. In this case we obtain an estimate which improves a recent result by L. Zhao [8].

Mathematics Subject Classification (2000): 11N35, 11L07, 11B57

Key words: large sieve, sparse set of moduli, square moduli, Farey fractions in short intervals, estimates on exponential sums

1 Introduction

Throughout this paper, we reserve the symbols c, c_i ($i = 1, 2, \dots$) for absolute constants. Further, we suppose that (a_n) is a sequence of complex numbers and that $Q, N \geq 1$. We set

$$(1) \quad S(\alpha) := \sum_{n \leq N} a_n e(n\alpha).$$

Bombieri's [1] classical large sieve inequality asserts that

$$(2) \quad \sum_{q \leq Q} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \ll (N + Q^2)Z,$$

where

$$Z := \sum_{n \leq N} |a_n|^2.$$

It is natural to ask whether (2) can be improved if the moduli q run over a sparse set $\mathcal{S}$ of natural numbers $\leq Q$. In the sequel, let S be the cardinality of $\mathcal{S}$.

P.D.T.A. Elliott [3] considered the case when $\mathcal{S}$ consists of primes. Essentially, he proved that

$$(3) \quad \sum_{p \in \mathcal{S}} \sum_{a=1}^{p-1} \left| S\left(\frac{a}{p}\right) \right|^2 \ll \left(\frac{N^2}{\log N} + QS \right) Z$$

and conjectured that the right-hand side can always be replaced by

$$(4) \quad \ll (N + QS)Z.$$

For the case when $\mathcal{S}$ consists of all primes $p \leq Q$ D. Wolke [7] found the estimate

$$(5) \quad \sum_{p \leq Q} \sum_{a=1}^{p-1} \left| S\left(\frac{a}{p}\right) \right|^2 \leq \frac{c}{1-\delta} \frac{Q^2 \log \log Q}{\log Q} Z$$

if $Q \geq 10$, $N = Q^{1+\delta}$, $0 < \delta < 1$.

L. Zhao [8] studied the case when $\mathcal{S}$ consists of squares. He established the bound

$$(6) \quad \sum_{q \leq Q_1} \sum_{\substack{a=1 \\ (a,q)=1}}^{q^2} \left| S\left(\frac{a}{q^2}\right) \right|^2 \ll (\log 2Q_1) \left(Q_1^3 + (N\sqrt{Q_1} + \sqrt{N}Q_1^2)N^\epsilon \right) Z.$$

The classical form (2) of the large sieve implies only the bound

$$(7) \quad \sum_{q \leq Q_1} \sum_{\substack{a=1 \\ (a,q)=1}}^{q^2} \left| S\left(\frac{a}{q^2}\right) \right|^2 \ll (N + Q_1^4)Z,$$

which is weaker than (6) if $Q_1 \gg N^{2/7+\epsilon}$. Using the later Lemma 1, which gives a general large sieve bound, one can also show that (see [8])

$$(8) \quad \sum_{q \leq Q_1} \sum_{\substack{a=1 \\ (a,q)=1}}^{q^2} \left| S\left(\frac{a}{q^2}\right) \right|^2 \ll Q_1(N + Q_1^2)Z,$$

which is weaker than (6) if $Q_1 \ll N^{1/2-\epsilon}$. So (6) is sharper than both (7) and (8) if $N^{2/7+\epsilon} \ll Q_1 \ll N^{1/2-\epsilon}$.

The aim of the present paper is to establish a general large sieve bound for the case when $\mathcal{S}$ is a sparse set which is in a certain sense well-distributed in arithmetic progressions. To do so, we shall use an extension of the method developed in [7]. Our main result, the later Theorem 2, will imply Wolke's estimate (5) and a sharpened version of Zhao's estimate (6). For the sake of its generality, the later Theorem 2 might have a number of other applications.

2 Statement of the results

In the sequel, for $t \in \mathbb{N}$ let

$$\mathcal{S}_t := \{q \in \mathbb{N} : tq \in \mathcal{S}\}$$

and $S_t := |\mathcal{S}_t|$. We first give a general estimate for the sum in question in terms of the number of elements of $\mathcal{S}_t$ in short segments of arithmetic progressions.

Theorem 1: *Suppose that $0 \leq M \leq Q$ and $\mathcal{S} \subset (M, M + Q]$. For $u > 0$, $t, k \in \mathbb{N}$, $l \in \mathbb{Z}$ define*

$$A_t(u, k, l) := \max_{M/t \leq y \leq (M+Q)/t} |\{q \in \mathcal{S}_t \cap (y, y + u] : q \equiv l \pmod{k}\}|.$$

Put

$$U := \begin{cases} 1, & \text{if } M < \sqrt{N}, \\ 0, & \text{otherwise.} \end{cases}$$

Then,

(9)

$$\sum_{q \in \mathcal{S}} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \leq c_1 N Z \left(U + \max_{r \leq \sqrt{N}} \max_{1/N \leq z \leq 1/(r\sqrt{N})} \max_{\substack{h \in \mathbb{Z} \\ (h,r)=1}} \right. \\ \left. \sum_{t|r} \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t)=1}} A_t\left(\frac{Q}{tzN}, \frac{r}{t}, hm\right) \right).$$

To simplify the estimate (9), we now set some natural conditions to the terms $A_t(u, k, l)$ with $(k, l) = 1$. We motivate these conditions by heuristic ideas.

Suppose that $t \leq \sqrt{N}$ and $0 < u \leq Q/t$. If the set $\mathcal{S}_t$ is nearly evenly distributed on the interval $(M/t, (M + Q)/t]$, we would expect that

$$A_t(u, 1, 0) \leq C \left(1 + \frac{S_t}{Q/t} \cdot u \right)$$

with $C \geq 1$ not too large. If we, more generally, assume the set $\mathcal{S}_t$ to be nearly evenly distributed in the residue classes mod k , we get

$$(10) \quad A_t(u, k, l) \leq C \left(1 + \frac{S_t/k}{Q/t} \cdot u \right).$$

For many sets $\mathcal{S}_t$ naturally appearing in arithmetic, the term on the right side of (10) will roughly give the correct order of magnitude, but in addition we have to take into consideration possible fluctuations on varying l . Therefore we multiply the right side of (10) by a non-negative constant $\delta_t(k, l)$ depending on the residue class $l \bmod k$. Thus, our first condition to $A_t(u, k, l)$ is

$$(11) \quad A_t(u, k, l) \leq C \left(1 + \frac{S_t/k}{Q/t} \cdot u \right) \delta_t(k, l).$$

Since $\sum_{l=1}^k A_t(u, k, l)$ should be roughly of the same size as $A_t(u, 1, 0)$, we may further assume that

$$(12) \quad \sum_{\substack{l=1 \\ (k, l)=1}}^k \delta_t(k, l) \leq k,$$

which is our second condition.

Finally, we suppose that

$$(13) \quad \delta_t(k, l) \leq X$$

for all $k \leq \sqrt{N}/t$, $l \in \mathbb{Z}$ with $(k, l) = 1$, where we think of X as a quantity which is small compared with Q and N .

In the sections 3 and 4 we will exemplify our heuristics by the cases when $\mathcal{S}$ consists of primes as well as of squares.

We shall later bound the right side of (9) by using the conditions (11), (12), (13). This shall lead us to the following

Theorem 2: *Suppose that $0 \leq M \leq Q$ and $\mathcal{S} \subset (M, M + Q]$. Define $A_t(u, k, l)$ and U as in Theorem 1. Assume the conditions (11), (12), (13) to hold for all $t \leq \sqrt{N}$, $k \leq \sqrt{N}/t$, $l \in \mathbb{Z}$ with $(k, l) = 1$, and $kQ/\sqrt{N} \leq u \leq Q/t$. Then,*

$$(14) \quad \sum_{q \in \mathcal{S}} \sum_{\substack{a=1 \\ (a, q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \leq c_2 C \left(NU + (\min\{QX, N\} + Q) \left(\sqrt{N} \log \log 10N + \max_{r \leq \sqrt{N}} \sum_{t|r} S_t \right) \right) Z.$$

Using $S_t \leq S$ for all $t \in \mathbb{N}$ and $\sum_{t|r} 1 \ll r^\varepsilon$, we deduce from Theorem 2

Corollary: *Let the conditions and assumptions of Theorem 2 be kept. Fix any $\varepsilon > 0$. Then,*

$$\sum_{q \in \mathcal{S}} \sum_{\substack{a=1 \\ (a, q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \ll \left(N + QN^\varepsilon X(\sqrt{N} + S) \right) Z.$$

This bound should be compared with Elliot's estimate (3) and conjecture (4) for the case when $\mathcal{S}$ consists of primes.

Employing Theorem 2 with $\mathcal{S}$ the set of squares lying in the interval $(Q, 2Q]$, we shall later derive the following improvement of Zhao's bound (6) from Theorem 2.

Theorem 3: *Suppose that $Q_1 \geq 1$. Fix any $\varepsilon > 0$. Then,*

$$(15) \quad \sum_{q \leq Q_1} \sum_{\substack{a=1 \\ (a, q)=1}}^{q^2} \left| S\left(\frac{a}{q^2}\right) \right|^2 \ll (\log \log 10NQ)^2 (Q_1^3 + N + N^{1/2+\varepsilon} Q_1^2) Z.$$

The bound (15) is sharper than the three bounds (6), (7) and (8) if $N^{1/4+\varepsilon} \ll Q_1 \ll N^{1/3-\varepsilon}$.

3 Theorem 2 implies Wolke's bound (5)

In this section we shall see that Theorem 2 contains Wolke's bound (5) as a special case.

As in [7], we assume that $Q \geq 10$, $N = Q^{1+\delta}$, $0 < \delta < 1$. Further, we choose $M := 0$ and $\mathcal{S}$ to be the set of all primes $\leq Q$.

First we consider the case when $t = 1$. Then, by the Brun-Titchmarsh inequality (see [5]), we have

$$(16) \quad A_1(u, k, l) \leq \frac{2u}{k \log u/k} \cdot \frac{k}{\varphi(k)}$$

if $u \geq k$.

For $u \geq kQ/\sqrt{N}$ the inequality (16), our assumption $N = Q^{1+\delta}$ and the prime number theorem yield

$$A_1(u, k, l) \leq \frac{4u}{(1-\delta)k \log Q} \cdot \frac{k}{\varphi(k)} \leq \frac{c_3 u S}{(1-\delta)kQ} \cdot \frac{k}{\varphi(k)}.$$

Thus, the condition (11) is satisfied if we set $C := c_3/(1-\delta)$ and $\delta_1(k, l) := k/\varphi(k)$. Obviously, the condition (12) is also satisfied, and the condition (13) holds with $X = c_4 \log \log Q$ (see [2]).

Now, we assume that $t \geq 2$. Then, $S_t \leq 1$. Thus, the conditions (11), (12), (13) hold trivially with $\delta_t(k, l) = 1$ and C, X chosen as above.

Therefore, we can apply Theorem 2 to this situation. Taking into account $S \leq c_5 Q / \log Q$ by the prime number theorem, we obtain the bound (5) from (14). $\square$

4 Proof of Theorem 3

Next, we shall derive Theorem 3 from Theorem 2.

If $Q_1^4 \leq N$, then (15) follows from (7). In the following, we assume that $Q_1^4 > N$.

First, we rewrite the sum in question in the form

$$(17) \quad \sum_{q \leq Q_1} \sum_{\substack{a=1 \\ (a,q)=1}}^{q^2} \left| S\left(\frac{a}{q^2}\right) \right|^2 = \sum_{q \in \mathcal{S}} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2,$$

where $\mathcal{S}$ is the set of squares $\leq Q_1^2$. We split up the set $\mathcal{S} \cap (\sqrt{N}, Q_1^2]$ into subsets of the form

$$\mathcal{S}(Q) := \mathcal{S} \cap (Q, 2Q],$$

where $Q \geq \sqrt{N}$. Our aim is to estimate the partial sums

$$\sum_{q \in \mathcal{S}(Q)} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2.$$

For the remaining sum we have

$$(18) \quad \sum_{q \in \mathcal{S} \cap (0, \sqrt{N}]} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \ll NZ$$

from (2).

As previously, we define

$$\mathcal{S}_t(Q) := \{q \in \mathbb{N} : tq \in \mathcal{S}(Q)\}$$

and $S_t(Q) := |\mathcal{S}_t(Q)|$. We now determine the set $\mathcal{S}_t(Q)$. Let $t = p_1^{v_1} \cdots p_n^{v_n}$ be the prime number factorization of t . For $i = 1, \dots, n$ let

$$u_i := \begin{cases} v_i, & \text{if } v_i \text{ is even,} \\ v_i + 1, & \text{if } v_i \text{ is odd.} \end{cases}$$

Put

$$f_t := p_1^{u_1/2} \cdots p_n^{u_n/2}.$$

Then $q = q_1^2 \in \mathcal{S}$ is divisible by t iff q_1 is divisible by f_t . Thus,

$$\mathcal{S}_t(Q) = \left\{ q_2^2 g_t : \sqrt{Q}/f_t < q_2 \leq \sqrt{2Q}/f_t \right\} \subset (Q/t, 2Q/t],$$

where

$$g_t := \frac{f_t^2}{t} = p_1^{u_1-v_1} \cdots p_n^{u_n-v_n}.$$

Hence,

$$(19) \quad S_t(Q) \leq \frac{\sqrt{2Q}}{f_t}.$$

As previously, we suppose that $k \in \mathbb{N}$, $l \in \mathbb{Z}$ and $(k, l) = 1$, and define

$$A_t(u, k, l) := \max_{Q/t \leq y \leq 2Q/t} |\{q \in \mathcal{S}_t(Q) \cap (y, y+u] : q \equiv l \pmod{k}\}|.$$

Let $\delta_t(k, l)$ be the number of solutions $x \pmod{k}$ to the congruence

$$(20) \quad x^2 g_t \equiv l \pmod{k}.$$

Then it is easily seen that the condition (11) with $S_t = S_t(Q)$ holds true for all positive $u \leq Q/t$ and some absolute constant $C \geq 1$. Clearly, (12) is also valid. The remaining task is to bound $\delta_t(k, l)$.

If $(g_t, k) > 1$, then $\delta_t(k, l) = 0$ since k and l are supposed to be coprime. Therefore, we can assume that $(g_t, k) = 1$. Let $g \pmod{k}$ be the multiplicative inverse of $g_t \pmod{k}$, i.e. $gg_t \equiv 1 \pmod{k}$. Put $l^* = gl$. Then (20) is equivalent to

$$(21) \quad x^2 \equiv l^* \pmod{k}.$$

Taking into account that $(k, l^*) = 1$, and using some elementary facts on the number of solutions of polynomial congruences modulo prime powers (see [6], for example), we see that (21) has at most 2 solutions if k is a power of an odd prime and at most 4 solutions if k is a power of 2. From this it follows that for all $k \in \mathbb{N}$ we have

$$\delta_t(k, l) \leq 2^{\omega(k)+1},$$

where $\omega(k)$ is the number of distinct prime divisors of k . For $k \leq \sqrt{N}$ we have

$$2^{\omega(k)} \ll N^\varepsilon$$

(see [2]). Therefore, (13) holds with

$$(22) \quad X := c_6 N^\varepsilon.$$

Now we can apply Theorem 2. Combining (14), (19) and (22), and taking into account that $Q \geq \sqrt{N}$, we obtain

$$(23) \quad \sum_{q \in \mathcal{S}(Q)} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \\ \leq c_7(\min\{QN^\varepsilon, N\} + Q) \left(\sqrt{N} \log \log 10N + \sqrt{Q} \max_{r \leq \sqrt{N}} \sum_{t|r} \frac{1}{f_t} \right) Z.$$

Next, we bound the function

$$\sigma(r) = \sum_{t|r} \frac{1}{f_t}.$$

Clearly, this function is multiplicative. If r is a prime power p^v , then

$$\sigma(r) \leq 1 + 2 \left(\frac{1}{p} + \frac{1}{p^2} + \dots \right) = 1 + \frac{2}{p-1} \leq \left(1 + \frac{1}{p-1} \right)^2 = \left(\frac{p^k}{\varphi(p^k)} \right)^2.$$

Hence, for all $r \in \mathbb{N}$ we have

$$\sigma(r) \leq \left(\frac{r}{\varphi(r)} \right)^2 \ll (\log \log 10r)^2.$$

Therefore, from (23), we obtain

$$(24) \quad \sum_{q \in \mathcal{S}(Q)} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \\ \leq c_8(\log \log 10NQ)^2 (\min\{QN^\varepsilon, N\} + Q) \left(\sqrt{N} + \sqrt{Q} \right) Z.$$

Considering the cases $QN^\varepsilon \leq N$, $N < QN^\varepsilon \leq N^{1+\varepsilon}$ and $N^{1+\varepsilon} < QN^\varepsilon$ separately, we see that the right-hand side of (24) is always bounded by

$$\leq c_9(\log \log 10NQ)^2 (Q^{3/2} + N^{1/2+\varepsilon}Q) Z =: T(Q).$$

Thus, we have

$$\sum_{q \in \mathcal{S} \cap (\sqrt{N}, Q_1^2]} \sum_{\substack{a=1 \\ (a,q)=1}}^q \left| S\left(\frac{a}{q}\right) \right|^2 \leq \sum_{j=0}^{\lceil \log_2(Q_1^2/\sqrt{N}) \rceil + 1} T\left(\frac{Q_1^2}{2^j}\right) \\ \ll (\log \log 10NQ_1)^2 (Q_1^3 + N^{1/2+\varepsilon} Q_1^2) Z.$$

Combining this with (18), we obtain (15). $\square$

5 Counting Farey fractions in short intervals

Our later proof of Theorem 1 relies on the following variant of the large sieve, which follows immediately from Theorem 2.11 in [4].

Lemma 1: *Let $(\alpha_r)_{r \in \mathbb{N}}$ be a sequence of real numbers and $(a_n)_{n \in \mathbb{N}}$ be a sequence of complex numbers. Define the trigonometrical polynomial $S(\alpha)$ as in (1). Suppose that $0 < \Delta \leq 1/2$ and $R \in \mathbb{N}$. Put*

$$K(\Delta) := \max_{\alpha \in \mathbb{R}} \sum_{\substack{r=1 \\ \|\alpha_r - \alpha\| \leq \Delta}}^R 1,$$

where $\|x\|$ denotes the distance of a real x to its closest integer. Then

$$\sum_{r=1}^R |S(\alpha_r)|^2 \leq c_{10} K(\Delta) (N + \Delta^{-1}) Z.$$

In our situation, the sequence $\alpha_1, \dots, \alpha_R$ equals the sequence of Farey fractions a/q with $q \in \mathcal{S}$, $1 \leq a \leq q$ and $(a, q) = 1$. For $\alpha \in \mathbb{R}$ put

$$I(\alpha) := [\alpha - \Delta, \alpha + \Delta] \quad \text{and} \quad P(\alpha) := \sum_{\substack{q \in \mathcal{S}, (a,q)=1 \\ a/q \in I(\alpha)}} 1.$$

Then we have

$$(25) \quad K(\Delta) = \max_{\alpha \in \mathbb{R}} P(\alpha).$$

The next lemma provides an estimate for $P(\alpha)$.

Lemma 2: *Suppose that $0 \leq M \leq Q$, $\mathcal{S} \subset (M, M + Q]$ and $0 < \Delta \leq 1/2$. Let $\alpha \in \mathbb{R}$. Define $A_t(u, k, l)$ as in Theorem 1. Put*

$$U := \begin{cases} 1, & \text{if } M < 1/\sqrt{\Delta}, \\ 0, & \text{otherwise.} \end{cases}$$

Then,

$$(26) \quad P(\alpha) \leq c_{11} \left(U + \max_{r \leq 1/\sqrt{\Delta}} \max_{\Delta \leq z \leq \sqrt{\Delta}/r} \max_{\substack{h \in \mathbb{Z} \\ (h, r) = 1}} \sum_{t|r} \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t) = 1}} A_t \left(\frac{\Delta Q}{tz}, \frac{r}{t}, hm \right) \right).$$

Proof: We adapt Wolke's method used to prove Lemma 1 in [7]. Let

$$(27) \quad \tau := \frac{1}{\sqrt{\Delta}}.$$

Then, by Dirichlet's approximation theorem, α can be written in the form

$$(28) \quad \alpha = \frac{b}{r} + z, \quad \text{where } r \leq \tau, (b, r) = 1, |z| \leq \frac{1}{r\tau}.$$

For $r \leq \tau$ we have

$$(29) \quad \frac{1}{r\tau} \geq \frac{1}{\tau^2} = \Delta.$$

We first note that we can restrict ourselves to the case when

$$(30) \quad z \geq \Delta.$$

If $|z| < \Delta$, then

$$P(\alpha) \leq P(b/r - \Delta) + P(b/r + \Delta).$$

Furthermore, by (29), we have $\Delta \leq 1/(r\tau)$. Therefore this case can be reduced to the case $|z| = \Delta$. Moreover, as $P(\alpha) = P(-\alpha)$, we can choose z positive. So we can assume (30).

Write

$$(31) \quad \Delta_1 := \frac{\Delta}{z} \quad (0 < \Delta_1 \leq 1),$$

$$(32) \quad j := \left\lceil \frac{1}{\Delta_1} \right\rceil + 1, \quad \Delta_2 := \frac{1}{j}.$$

Then, obviously,

$$(33) \quad \frac{\Delta_1}{2} \leq \Delta_2 \leq \Delta_1.$$

Put

$$(34) \quad y_i := M + i\Delta_2 Q, \quad Y_i = (y_i, y_{i+1}], \quad (i = 0, 1, \dots)$$

and

$$(35) \quad P(\alpha) = \sum_{i=0}^{j-1} \sum_{q \in \mathcal{S} \cap Y_i} \sum_{\substack{(a,q)=1 \\ a/q \in I(\alpha)}} 1 =: \sum_{i=0}^{j-1} P_i(\alpha), \quad \text{say.}$$

Now, for $0 \leq i \leq j-1$ and

$$(36) \quad q \in Y_i, \quad (a, q) = 1, \quad \frac{a}{q} \in I(\alpha),$$

we have $q(\alpha - \Delta) \leq a \leq q(\alpha + \Delta)$ or, by (28) and (30), $y_i r(z - \Delta) \leq ar - bq \leq y_{i+1} r(z + \Delta)$. Write

$$(37) \quad W_i := [w_i, w'_i] := [y_i r(z - \Delta), y_{i+1} r(z + \Delta)].$$

Thus, (36) implies $ar - bq = m$, $m \in W_i$. If $m = 0$, then $q = r$ since $(a, q) = 1 = (b, r)$. Hence,

$$(38) \quad P_i(\alpha) \leq \nu_i + \sum_{\substack{m \in W_i \\ m \neq 0}} \sum_{\substack{q \in \mathcal{S} \cap Y_i \\ q \equiv -\bar{b}m \pmod{r}}} 1,$$

where

$$\nu_i := \begin{cases} 1, & \text{if } r \in Y_i, \\ 0, & \text{otherwise,} \end{cases}$$

and $\bar{b} \bmod r$ is the multiplicative inverse of $b \bmod r$, *i.e.* $\bar{b}b \equiv 1 \bmod r$.

For $t \in \mathbb{N}$ write

$$W_i/t := [w_i/t, w'_i/t], \quad Y_i/t := (y_i/t, y_{i+1}/t].$$

Then from (38) it follows that

$$(39) \quad P_i(\alpha) \leq \nu_i + \sum_{t|r} \sum_{\substack{m \in W_i/t \\ (m, r/t) = 1 \\ m \neq 0}} \sum_{\substack{q \in \mathcal{S}_t \cap (Y_i/t) \\ q \equiv -\bar{b}m \bmod r/t}} 1.$$

By (31), (33) and (34), we get $y_{i+1} - y_i \leq \Delta Q/z$. Thus, from (39), we obtain

$$(40) \quad P_i(\alpha) \leq \nu_i + \sum_{t|r} \sum_{\substack{m \in W_i/t \\ (m, r/t) = 1 \\ m \neq 0}} A_t \left(\frac{\Delta Q}{tz}, \frac{r}{t}, -\bar{b}m \right).$$

From (31), (32), (33), (34) and (37) it follows that $w'_i \leq rz(y_{i+1} + 2Q\Delta_2) = rzy_{i+3}$ and, similarly, $w_i \geq rzy_{i-2}$. Moreover, we have $w_0 \geq 0$ by (30) and $M \geq 0$. Thus, the intervals W_i ($0 \leq i \leq j-1$) cover the interval $V = [0, rzy_{j+2}]$ at most 6 times. Furthermore, by (32), (34) and $M \leq Q$, we have $rzy_{j+2} \leq 4rzQ$. Hence, (35) and (40) imply

$$(41) \quad P(\alpha) \leq U + 6 \sum_{t|r} \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t) = 1}} A_t \left(\frac{\Delta Q}{tz}, \frac{r}{t}, -\bar{b}m \right),$$

where U is defined as in Lemma 2. From (27), (28), (30) and (41), we deduce (26). This completes the proof. $\square$

6 Proofs of Theorems 1 and 2

We are now in a position to prove our Theorems 1 and 2.

Proof of Theorem 1: Taking $\Delta := 1/N$, the result of Theorem 1 follows from Lemma 1, (25) and Lemma 2. $\square$

Proof of Theorem 2: Suppose that $r \leq \sqrt{N}$, $1/N \leq z \leq 1/(r\sqrt{N})$, $(h, r) = 1$ and $t|r$. Then, by condition (11), we have

$$(42) \quad \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t) = 1}} A_t \left(\frac{Q}{tzN}, \frac{r}{t}, hm \right) \leq C \left(1 + \frac{tS_t}{rzN} \right) F_h(r, t),$$

where

$$F_h(r, t) := \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t) = 1}} \delta_t(r/t, hm).$$

If $z \geq 1/(4Q)$, then the condition (12) gives

$$(43) \quad F_h(r, t) \leq \frac{8rzQ}{t}.$$

If $z < 1/(4Q)$, then the conditions (12) and (13) imply

$$(44) \quad F_h(r, t) \leq \min \left\{ \frac{4rzQX}{t}, \frac{r}{t} \right\}.$$

For $r \leq \sqrt{N}$ and $1/N \leq z \leq 1/(r\sqrt{N})$, from (42), (43) and (44), we derive

$$(45) \quad \begin{aligned} & N \sum_{t|r} \sum_{\substack{0 < m \leq 4rzQ/t \\ (m, r/t) = 1}} A_t \left(\frac{Q}{tzN}, \frac{r}{t}, hm \right) \\ & \leq c_{12} C (\min \{QX, N\} + Q) \left(\sqrt{N} \sum_{t|r} \frac{1}{t} + \sum_{t|r} S_t \right) \end{aligned}$$

by a short calculation. From Theorem 1, (45) and

$$\sum_{t|r} \frac{1}{t} \leq \prod_{p|r} \left(1 + \frac{1}{p} + \frac{1}{p^2} + \dots \right) = \prod_{p|r} \frac{p}{p-1} = \frac{r}{\varphi(r)} \leq c_{13} \log \log 10r,$$

we obtain the result of Theorem 2. $\square$

Acknowledgement. This paper was written when the author held a post-doctoral position at the Harish-Chandra Research Institute at Allahabad (India). The author wishes to thank this institute for financial support.

References

- [1] Bombieri, E., *On the large sieve*, Mathematika 12 (1965) 201-225.
- [2] B. Crstici, D.S. Mitrinović, J. Sándor, *Handbook of number theory*, Kluwer Academic Publishers Group, Dordrecht, 1996.
- [3] P.D.T.A. Elliott, *On inequalities of large sieve type*, Acta Arith. 18 (1971) 405-422.
- [4] H.L. Montgomery, *Topics in multiplicative number theory*, Lecture Notes in Mathematics, Vol. 227, Springer-Verlag, Berlin-New York, 1971.
- [5] H.L. Montgomery, R.C. Vaughan, *The large sieve*, Mathematika 20 (1973) 119-134.
- [6] I. Niven, H.S. Zuckerman, H.L. Montgomery, *An introduction to the theory of numbers*, John Wiley & Sons, New York, 1991.
- [7] D. Wolke, *On the large sieve with primes*, Acta Math. Acad. Sci. Hungar. 22 (1971/72) 239-247.
- [8] L. Zhao, *Large sieve inequality with characters to square moduli*, Acta Arith. 112 (2004) 297-308.